



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 062 879 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
27.12.2000 Bulletin 2000/52

(51) Int Cl.7: **A23L 1/305, A23L 1/304,
A61K 31/195, A61K 33/00**

(21) Application number: **99112319.1**

(22) Date of filing: **26.06.1999**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(74) Representative:
**Jönsson, Hans-Peter, Dr.Dipl.-Chem. et al
Patentanwälte
von Kreisler Selting Werner
Postfach 10 22 41
50462 Köln (DE)**

(71) Applicant: **B. BRAUN MELSUNGEN AG
34212 Melsungen (DE)**

(72) Inventors:
• **Guardiola, Jaime Dr.
66424 Homburg (DE)**
• **Wolf, Martin Dr.
34212 Melsungen (DE)**

Remarks:

A request for correction dated 05.08.99 has been filed pursuant to Rule 88 EPC. A decision on the request will be taken during the proceedings before the Examining Division (Guidelines for Examination in the EPO, A-V, 3.).

(54) **Aqueous solution for the parenteral nutrition**

(57) The invention refers to an aqueous solution for the parenteral nutrition of mammals consisting of:

a) a pattern of amino acids calculated on the basis of 100 g/l of amino acids:

Isoleucine	4.0 - 5.5 g/l
Leucine	8.0 - 10.0 g/l
Lysine	6.0 - 8.0 g/l
Methionine	4.0 - 6.0 g/l
Phenylalanine	4.0 - 6.0 g/l
Threonine	4.0 - 6.0 g/l
Tryptophane	1.0 - 2.0 g/l
Valine	6.0 - 8.0 g/l
Arginine	10.0 - 12.0 g/l
Histidine	1.5 - 3.5 g/l
Alanine	9.0 - 12.0 g/l

(continued)

Aminoacetic Acid (Glycine)	11.0 - 16.0 g/l
Asparagine	0 - 1.0 g/l
Aspartic Acid	5.5 - 8.0 g/l
Acetylcysteine	0 - 2.5 g/l
Glutamic Acid	6.0 - 10.0 g/l
Ornithine	0 - 1.0 g/l
Proline	4.0 - 6.0 g/l
Serine	1.0 - 3.0 g/l
Tyrosine*	0.1 - 0.5 g/l
*as Acetyltyrosine	0 - 2.0 g/l
Taurine	0 - 4.0 g/l

EP 1 062 879 A1

b) a pattern of electrolytes of:

Sodium	0 - 100.0 mmol/l
Potassium	0 - 80.0 mmol/l
Magnesium	0 - 8.0 mmol/l
Calcium	0 - 8.0 mmol/l and/or
Zinc	0 - 0.08 mmol/l
Phosphate	0 - 20.0 mmol/l
Chloride	0 - 120.0 mmol/l
Acetate	0 - 120.0 mmol/l

(continued)

Citric acid	0 - 10.0 mmol/l
Malate	0 - 80.0 mmol/l
Lactate	0 - 10.0 mmol/l
Glycerophosphate	0 - 30.0 mmol/l and/or
Gluconate	0 - 120.0 mmol/l

and

c) optionally a carbohydrate solution and/or a fat emulsion.

Description

[0001] The invention refers to an aqueous solution for the parenteral nutrition of mammals.

[0002] EP 0 671 166 A2 discloses an infusion preparation for nutrient supply use. It comprises sugar, amino acids, electrolytes and fat emulsion. The examples disclose several patterns of amino acids.

[0003] In the same way DE 43 16 326 C1 discloses different patterns of amino acids for the total parenteral nutrition by perivenous application.

[0004] Similar to that disclosure of DE 43 16 326 C1, DE 39 16 903 A1 discloses an aqueous composition of parenteral nutrition which is in particular defined by a specific amino acid pattern.

[0005] R. Zander, Infusionsther. Transfusionsmed. 1993;20:217 - 235 discloses a specific base concept for infusion solutions. In particular, the concepts of base excess (BE mmol/l and BE pot. mmol/l) should be introduced, in order to give an indication of potential alterations in the HCO_3^- pool after infusion and metabolism of infusion solutions. This also applies to blood derivatives, where transfusion acidosis as well as alkalosis may occur, and in hemodialysis and peritoneal dialysis where the occurrence of acidosis and alkalosis during therapy need to be reckoned with. The above mentioned patterns of amino acid solutions partially available in the market do not fulfill the requirements of the concept of a base excess.

[0006] Thus, it was the object of the invention to provide an aqueous solution for the parenteral nutrition of mammals fulfilling the requirements of said concept.

[0007] The solution of the above mentioned object is fulfilled in a first embodiment by an aqueous solution of the parenteral nutrition of mammals consisting of

a) a pattern of amino acids calculated on the basis of 100 g/l of amino acids

Isoleucine	4.0 - 5.5 g/l
Leucine	8.0 - 10.0 g/l
Lysine	6.0 - 8.0 g/l
Methionine	4.0 - 6.0 g/l
Phenylalanine	4.0 - 6.0 g/l
Threonine	4.0 - 6.0 g/l
Tryptophane	1.0 - 2.0 g/l
Valine	6.0 - 8.0 g/l
Arginine	10.0 - 12.0 g/l
Histidine	1.5 - 3.5 g/l
Alanine	9.0 - 12.0 g/l
Aminoacetic Acid (Glycine)	11.0 - 16.0 g/l
Asparagine	0 - 1.0 g/l
Aspartic Acid	5.5 - 8.0 g/l
Acetylcysteine	0 - 2.5 g/l
Glutamic Acid	6.0 - 10.0 g/l
Ornithine	0 - 1.0 g/l
Proline	4.0 - 6.0 g/l
Serine	1.0 - 3.0 g/l
Tyrosine*	0.1 - 0.5 g/l
*as Acetyltyrosine	0 - 2.0 g/l
Taurine	0 - 4 g/l

b) a pattern of electrolytes of

Sodium	0 - 100.0 mmol/l
Potassium	0 - 80.0 mmol/l
Magnesium	0 - 8.0 mmol/l
Calcium	0 - 8.0 mmol/l and/or
Zinc	0 - 0.08 mmol/l
Phosphate	0 - 20.0 mmol/l

(continued)

Chloride	0 - 120.0 mmol/l
Acetate	0 - 120.0 mmol/l
Citric acid	0 - 10.0 mmol/l
Malate	0 - 80.0 mmol/l
Lactate	0 - 10.0 mmol/l
Glycerophosphate	0 - 30.0 mmol/l and/or
Gluconate	0 - 120.0 mmol/l

c) optionally a carbohydrate solution (sugar) and/or fat emulsion.

[0008] Parenteral nutrition must supply the body with all the components necessary for growth and tissue generation. The amino acids play a prominent role since they are the basic components for protein synthesis. In order ensure optimal utilisation of the amino acids, the administration of an additional energy source is required. This can be fulfilled partly in the form of the optionally used carbohydrates. As glucose can be utilized directly, it is the carbohydrate of choice, but in the same way Xylitol can be utilized instead of or partially substituting glucose.

[0009] However, the prior art of course discloses the use of other carbohydrates.

[0010] Electrolytes are administered for the maintenance of metabolic and physiologic function. They ensure an adequate parenteral nutritional supply with the aqueous solution.

[0011] Following intravenous infusion the constituents of the solution according to the present invention are immediately available for metabolism due to the specific pattern of the ingredients. Therefore, no further investigations on the bioavailability are necessary. A portion of the amino acids is used for proteins synthesis, the rest being broken as following: The amino groups are separated by transamination and the carbon moiety is either oxidized to CO₂ in the citrate cycle or utilised in the liver as a substrate for a gluconeogenesis. The amino groups resulting for protein breakdown in muscle tissue are transported to the liver, where urea is synthesized. The ingredients of the aqueous solution according to the present invention are naturally occurring substances or metabolic intermediates and their concentration after appropriate infusion is similar to physiological levels.

[0012] The overall content of the amino acids may be varied according to the necessity of the patient to be administered therewith. Thus, the overall content of the amino acids in aqueous solutions according to the present invention may vary between 20 to 180 g/l. Nevertheless, it is to be ensured that the scope of pattern as mentioned above is not left. Accordingly, the given values have to be divided or multiplied accordingly for higher or lower concentration.

[0013] In a further preferred embodiment of the present invention, the pH value of the solutions has been optimized regarding physiologic acid-base-balance. Accordingly, it is in particular preferred that the pH values of the aqueous solutions are within the range of 4.8 to 6.5.

[0014] The use of at least 20 % by weight of branched amino acids is in particular preferred. W. Hartig, Moderne Infusionstherapie, Parenterale Ernährung, VEB Johann Ambrosius Barth 1989, pages 451 to 455 describes a preferred selection of amino acids according to Rose. The table on page 451 relating to essential amino acids only indicates an amount of the branched amino acids isoleucine, leucine and valine in an amount of 40,9 %. Corresponding to a preferred amino acid solution having a content of essential amino acids in an amount of 42 %. This amount of branched amino acids corresponds to an amount of 17,2 % by weight.

[0015] On the other hand, according to the present invention, a significant enlargement of the amount of branched amino acids was available in comparison to the above Rose pattern (approximately 18 % by weight). This improvement leads to an approximation to the amino acid pattern of egg protein for which the complete bioavailability is assumed.

[0016] In this respect not only the amount of branched amino acids is of importance but the relation to the amount in relation other amino acids. In particular the branched amino acids dominate (in particular limit) the solubility of the amino acids as such due to the relative pure solubility thereof. According to the knowledge of the inventors there is no other amino acid solution known or available which has an amino acid concentration of at least 15 % by weight containing an amount of branched amino acids of 20 % by weight based on the total amount of amino acids or more. Accordingly, the amount of branched amino acids is in particular of importance for amino acid solutions of the high preferable concentration such as 16 or 18 % by weight of amino acids.

[0017] It has also been considered that the potential base excess of the used amino acid solution does not exceed 20 mmol/day by using the maximum recommended dosage of amino acids 2 g protein/kg body weight and day.

[0018] By choosing the electrolyte composition (chloride, acetate, malate, citrate in the case of the electrolyte containing solutions) as well as by using the amino acid salts alone (lysine/lysine acetate but also the available aspartates, glutamates and pyro-glutamates) it had been considered that the potential base excess should be preferably between -10 and +10 mmol/l in the final product. For the electrolyte containing variations of the aqueous solution the potential

base excess should be preferably close to 0 (+/- 5 mmol/l). For the electrolyte-free variations of the solutions according to the present invention the possibility of the base excess adjustment is limited because the solutions have to be chloride-free and therefore, the potential base excess varies preferably in the range of -15 to -8 mmol/l in these solutions. The relatively negative base excess of the electrolyte-free variations of the solutions according to the present invention causes no problems, if in particular the electrolyte-free amino acid solution is part of the therapy conception.

[0019] Electrolyte-free amino acid solutions must be mixed with electrolyte solutions to balance the electrolyte intake of the patient during the administration. The compositions according to the present invention are in particular based on the experience in the field of known amino acid solutions which are in the market for many years. The specific pattern of the amino acids according to the present invention is the result of many years of clinical experiments with solutions available in the market, now with the new development taking into account the physiological aspect of the base excess, the galenic aspect of the formulations and the stability of the product in addition to the amino acid pattern. In view of the high solubility of the amino acids when following the above mentioned pattern a great variation of the amount of amino acid and electrolyte can be provided. Thus, the same base solution may be the basis of a 10 % solution (corresponding to 100 g/l amino acids to correspond to other 3, 5, 8 or even 16 to 18 % concentrations).

[0020] In case the aqueous solution according to the present invention contains amino acids and carbohydrates like glucose in one container, usually a reaction between these components take place depending mainly on the temperature and also on oxygen content of the mixture. The so-called Maillard-reaction causes a yellow color of the final product. This is in particular in the case of total parenteral solutions being in the market. These solutions are therefore produced aseptically without the usual heat sterilization at temperatures above 121 °C.

[0021] The reaction between glucose and amino acids ends up with the formation of Maillard-products (yellow/brown colored condensation products), thus resulting in a concentration loss of some of the available amino acids in the solution. This phenomenon has been described extensively in the literature. It is important to know in this connection the influence of the physiological salts (phosphate, glycerophosphate, acetate, chloride, citrate, lactate and malate) on the reaction kinetic of the above-mentioned Maillard-reaction, influencing lately the stability of the solution and finally the shelflife of the medical preparation. This connection is not or only rarely investigated and published. The various solutions according to the present invention are optimized regarding the base excess as well as the kinetic of the above mentioned Maillard-reaction.

[0022] The base excess may be calculated as following:

$$\text{Base excess (BE)} = \text{TA} + 24$$

TA: Titration acidity (mmol/l NaOH for reaching pH value 7.4)

24 mmol/l Na bicarbonate in blood

$$\text{Potential BE (Pot. BE)} = \text{ME} + \text{BE}$$

ME: Metabolic effect (metabolization of R-COOH > CO₂)

e.g.: ME = (Acetate-Concentration) + (Lactate-Concentration) + 2 (Malate-Concentration) + 3 (Citrate-Concentration)

[0023] In a further preferred embodiment the aqueous solution according to the present invention are characterized in that the pattern of the amino acids calculated on the basis of 100 g/l of the amino acids is:

Isoleucine	4.7 - 5.3 g/l
Leucine	8.5 - 9.5 g/l
Lysine	6.5 - 7.5 g/l
Methionine	4.0 - 5.0 g/l
Phenylalanine	4.5 - 5.5 g/l
Threonine	4.0 - 5.0 g/l
Tryptophane	1.5 - 1.7 g/l
Valine	6.0 - 7.0 g/l
Arginine	11.0 - 12.0 g/l
Histidine	2.5 - 3.5 g/l
Alanine	10.0 - 11.0 g/l

(continued)

Aminoacetic Acid (Glycine)	11.0 - 12.0 g/l
Asparagine	0 - 0.5 g/l
Aspartic Acid	5.5 - 6.0 g/l
Acetylcysteine	0.2 - 1.0 g/l**
Glutamic Acid	6.0 - 8.0 g/l
Ornithine	0 - 0.5 g/l
Proline	5.0 - 6.0 g/l
Serine	2.0 - 3.0 g/l
Tyrosine*	0.3 - 0.5 g/l
*as Acetyltyrosine	0 - 20 g/l
Taurine	0 - 2.5 g/l

** susceptible to oxidation, thus, dosage may depend on manufacturing technology and container as well

[0024] In the same way a further embodiment of the present invention is characterized in that the pattern of electrolytes being:

Sodium	25.0 - 75.0 mmol/l
Potassium	20.0 - 30.0 mmol/l
Magnesium	20.0 - 30.0 mmol/l
Calcium	0 - 4.0 mmol/l
Zinc	0 - 0.04 mmol/l
Phosphate	5.0 - 15.0 mmol/l
Chloride*	25.0 - 75.0 mmol/l
Acetate*	25.0 - 75.0 mmol/l
Citric acid	1.0 - 3.0 mmol/l
Malate	0 - 40.0 mmol/l
Lactate	0 - 5.0 mmol/l
Glycerophosphate	5 - 30.0 mmol/l and/or
Gluconate	25 - 120.0 mmol/l

[0025] The expression of the term "mammals" according to the present invention of course includes all kinds of mammals, in particular human beings.

Embodiment:

[0026]

Amino Acid %	100 [g/l]
Isoleucine	5.00
Leucine	8.90
Lysine	6.81
Methionine	4.40
Phenylalanine	4.70
Threonine	4.20
Tryptophane	1.60
Valine	6.20
Arginine	11.50
Histidine	3.00
Alanine	10.50
Aminoacetic Acid (Glycine)	12.00
Asparagine	

(continued)

Amino Acid %	100 [g/l]
Aspartic Acid	5.60
Acetylcysteine	0.21
Glutamic Acid	7.20
Ornithine	
Proline	5.50
Serine	2.30
Tyrosine	0.40

Electrolytes	10 [mol/l]
Sodium	50.0
Potassium	25.0
Magnesium	2.5
Calcium	-
Phosphate	10.0
Zinc	-
Chloride	52.0
Acetate	46.0
Citric acid	2.0
Malate	-
Lactate	-
Glycerophosphate	-
Gluconate	-

Claims

1. Aqueous solution of the parenteral nutrition of mammals consisting of:

a) a pattern of amino acids calculated on the basis of 100 g/l of amino acids:

Isoleucine	4.0 - 5.5 g/l
Leucine	8.0 - 10.0 g/l
Lysine	6.0 - 8.0 g/l
Methionine	4.0 - 6.0 g/l
Phenylalanine	4.0 - 6.0 g/l
Threonine	4.0 - 6.0 g/l
Tryptophane	1.0 - 2.0 g/l
Valine	6.0 - 8.0 g/l
Arginine	10.0 - 12.0 g/l
Histidine	1.5 - 3.5 g/l
Alanine	9.0 - 12.0 g/l
Aminoacetic Acid (Glycine)	11.0 - 16.0 g/l
Asparagine	0 - 1.0 g/l
Aspartic Acid	5.5 - 8.0 g/l
Acetylcysteine	0 - 2.5 g/l
Glutamic Acid	6.0 - 10.0 g/l
Ornithine	0 - 1.0 g/l
Proline	4.0 - 6.0 g/l
Serine	1.0 - 3.0 g/l

(continued)

Tyrosine*	0.1 - 0.5 g/l
*as Acetyltyrosine	0 - 2.0 g/l
Taurine	0 - 4.0 g/l

b) a pattern of electrolytes of:

Sodium	0 - 100.0 mmol/l
Potassium	0 - 80.0 mmol/l
Magnesium	0 - 8.0 mmol/l
Calcium	0 - 8.0 mmol/l and/or
Zinc	0 - 0.08 mmol/l
Phosphate	0 - 20.0 mmol/l
Chloride	0 - 120.0 mmol/l
Acetate	0 - 120.0 mmol/l
Citric acid	0 - 10.0 mmol/l
Malate	0 - 80.0 mmol/l
Lactate	0 - 10.0 mmol/l
Glycerophosphate	0 - 30.0 mmol/l and/or
Gluconate	0 - 120.0 mmol/l

and

c) optionally a carbohydrate solution and/or a fat emulsion.

2. Solution according to claim 1, characterized in that the content of amino acid varies in the range of 20 to 180 g/l.
3. Solution according to claim 1, characterized in that the pH value is in the range of 4.8 to 6.5.
4. Solution according to claim 1, characterized by an amount of 20 % by weight of branched amino acids.
5. Solution according to claim 1, characterized by an amount of at least 42 % by weight of essential amino acids.
6. Solution according to claim 1, characterized by an amount of at least 24 % of precursors selected from Asp, Glu and/or Arg.
7. Solution according to claim 1, characterized in that the potential base excess is within the range of -10 to + 10 mmol/l.
8. Solution according to claim 1, characterized in being sulfite-free.
9. Solution according to claim 1, characterized in that the pattern of the amino acids calculated on the basis of 100 g/l of amino acids is:

Isoleucine	4.7 - 5.3 g/l
Leucine	8.5 - 9.5 g/l
Lysine	6.5 - 7.5 g/l
Methionine	4.0 - 5.0 g/l
Phenylalanine	4.5 - 5.5 g/l
Threonine	4.0 - 5.0 g/l
Tryptophane	1.5 - 1.7 g/l
Valine	6.0 - 7.0 g/l
Arginine	11.0 - 12.0 g/l
Histidine	2.5 - 3.5 g/l

(continued)

Alanine	10.0 - 11.0 g/l
Aminoacetic Acid (Glycine)	11.0 - 12.0 g/l
Asparagine	0 - 0.5 g/l
Aspartic Acid	5.5 - 6.0 g/l
Acetylcysteine	0.2 - 1.0 g/l
Glutamic Acid	6.0 - 8.0 g/l
Ornithine	0 - 0.5 g/l
Proline	5.0 - 6.0 g/l
Serine	2.0 - 3.0 g/l
Tyrosine*	0.1 - 0.5 g/l
*as Acetyltyrosine	0 - 0.1 g/l
Taurine	0 - 2.5 g/l

10. Solution according to claim 1, characterized in that the pattern of electrolytes being:

Sodium	25,0 - 75,0 mmol/l
Potassium	20,0 - 30,0 mmol/l
Magnesium	20,0 - 30,0 mmol/l
Calcium	0 - 4.0 mmol/l and/or
Zinc	0 - 0.04 mmol/l
Phosphate	5.0 - 15.0 mmol/l
Chloride	25.0 - 75.0 mmol/l
Acetate	25.0 - 75.0 mmol/l
Citric acid	1.0 - 3.0 mmol/l
Malate	0 - 40.0 mmol/l
Lactate	0 - 5.0 mmol/l
Glycerophosphate	5 - 30.0 mmol/l and/or
Guconate	25 - 120.0 mmol/l



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 11 2319

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	EP 0 712 583 A (TANABE SEIYAKU) 22 May 1996 (1996-05-22) * claims; tables 1-12 * ---	1-10	A23L1/305 A23L1/304 A61K31/195 A61K33/00
A	EP 0 705 542 A (SANDOZ NUTRITION) 10 April 1996 (1996-04-10) * claims * ---	1-10	
A	US 5 206 269 A (J.K.LUDWIG ET AL.) 27 April 1993 (1993-04-27) * claims * ---	1-10	
D,A	DE 43 16 326 C (FRESENIUS) 9 June 1994 (1994-06-09) * the whole document * ---	1-10	
D,A	DE 39 16 903 A (LEOPOLD PHARMA) 28 February 1991 (1991-02-28) * the whole document * ---	1-10	
D,A	EP 0 671 166 A (GREEN CROSS) 13 September 1995 (1995-09-13) * claims; examples * -----	1-10	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A23L A61K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 16 December 1999	Examiner Van Moer, A
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03 82 (Pac01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 11 2319

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

16-12-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 712583 A	22-05-1996	JP 8143459 A	04-06-1996
		CN 1129104 A	21-08-1996
		SG 34308 A	06-12-1996
		US 5767123 A	16-06-1998
EP 705542 A	10-04-1996	US 5719133 A	17-02-1998
		AT 161397 T	15-01-1998
		CA 2158635 A,C	22-03-1996
		DE 69501306 D	05-02-1998
		DE 69501306 T	18-06-1998
		ES 2111384 T	01-03-1998
		GR 3026072 T	29-05-1998
		HK 1005768 A	22-01-1999
		SG 33502 A	18-10-1996
		SI 705542 T	30-04-1998
		US 5719134 A	17-02-1998
US 5206269 A	27-04-1993	AU 3371593 A	23-09-1993
		CA 2091602 A	21-09-1993
		EP 0561141 A	22-09-1993
		JP 6024971 A	01-02-1994
DE 4316326 C	09-06-1994	AT 171847 T	15-10-1998
		DE 59407031 D	12-11-1998
		EP 0625313 A	23-11-1994
		ES 2124810 T	16-02-1999
		FI 942216 A	16-11-1994
		NO 941506 A	16-11-1994
DE 3916903 A	28-02-1991	AT 88065 T	15-04-1993
		DK 399341 T	24-05-1993
		EP 0399341 A	28-11-1990
EP 671166 A	13-09-1995	JP 5031151 A	09-02-1993
		JP 5032540 A	09-02-1993
		JP 5032541 A	09-02-1993
		JP 2950348 B	20-09-1999
		JP 5009111 A	19-01-1993
		JP 5065220 A	19-03-1993
		EP 0752243 A	08-01-1997
		CA 2067062 A	27-10-1992
		EP 0510687 A	28-10-1992
		JP 5301825 A	16-11-1993
		US 5972367 A	26-10-1999
		US 5674527 A	07-10-1997
		US 5626880 A	06-05-1993

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 99 11 2319

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

16-12-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 671166 A		JP 5148149 A	15-06-1993

EPO FORM P0455

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82